

Antimicrobial susceptibility results

1/1/2019-11/1/2019

Gram positive cocci	<i>Enterococcus faecalis</i>	<i>Enterococcus faecium</i>	<i>Staphylococcus aureus</i>	MSSA	MRSA	<i>Staphylococcus epidermidis</i>	<i>Staphylococcus lugdunensis</i>	Group B <i>Streptococcus</i>	<i>Streptococcus pneumoniae</i>
Max strains tested	231	38	1,267	1,006	330	92	81	54	41
% Susceptible									
Amoxicillin/clavulanate	-	-	-	-	0	-	-	-	95
Ampicillin	99	30	-	-	0	-	-	-	-
Ciprofloxacin	80*	24*	78	93	36	72	99	-	-
Clindamycin	-	-	76	77	69	65	76	61	93
Erythromycin	-	-	54**	66**	16**	36**	76**	-	61
Daptomycin	100	-	100	100	100	100	100	-	-
Linezolid	99	100	99	100	99	100	-	-	100
Oxacillin/Nafcillin	-	-	74	100	0	38	90	-	-
Tetracycline	29	-	94***	94***	93***	78***	97***	-	76
Trimethoprim/sulfa	-	-	96	96	96	55	96	-	-
Vancomycin	97	45	100	100	100	100	100	-	100
Penicillin (CSF)	-	-	-	-	-	-	-	-	73
Penicillin (Non-CSF)	-	-	-	-	-	-	-	-	100
Ceftriaxone (CSF)	-	-	-	-	-	-	-	-	95
Ceftriaxone (Non-CSF)	-	-	-	-	-	-	-	-	98
Cefazolin	-	-	74	100	0	34	89	-	-
Ceftaroline	-	-	100	-	100	-	-	-	-
For urinary tract infections only									
Nitrofurantoin	99	8	99	99	99	100	100	-	-

*Should only be used to treat urinary tract infections caused by *Enterococcus* species.

**Susceptibility to erythromycin can be used to infer susceptibility to azithromycin.

***Susceptibility to tetracycline can be used to infer susceptibility to doxycycline.